






Glueballs, Hybrids & Exotics

An Experimental & Phenomenological Overview




Curtis A. Meyer
Carnegie Mellon University
May 31, 2006





Outline of Talk

- Introduction
- Meson Spectroscopy
- Glueballs
 - Expectations
 - Experimental Data
 - Interpretation
- Hybrid Mesons
 - Expectations
 - Experimental Data
- Summary and Future




Normal Mesons

Non-quark-antiquark

0⁻⁻ 0⁺⁻ 1⁻⁺ 2⁺⁻ 3⁻⁺ ...

quark-antiquark pairs



↑ orbital

L=2

$\rho_3, \omega_3, \phi_3, K_3$	3 ⁻⁻
$\rho_2, \omega_2, \phi_2, K_2$	2 ⁻⁻
$\rho_1, \omega_1, \phi_1, K_1$	1 ⁻⁻
$\pi_2, \eta_2, \eta'_2, K_2$	2 ⁻⁺

radial



L=1

a_2, f_2, f'_2, K_2	2 ⁺⁺
a_1, f_1, f'_1, K_1	1 ⁺⁺
a_0, f_0, f'_0, K_0	0 ⁺⁺
b_1, h_1, h'_1, K_1	1 ⁺⁻

L=0


ρ, ω, ϕ, K^*	1 ⁻⁻
π, η, η', K	0 ⁻⁺



$$J=L+S \quad (2S+1) L_J$$

$$P=(-1)^{L+1}$$


$$C=(-1)^{L+S} \quad {}^1S_0 = 0^{-+}$$

$$G=C(-1)^I \quad {}^3S_1 = 1^{--}$$





Spectrum




QCD is a theory of quarks and gluons


What role do gluons play in the meson spectrum?

Lattice calculations predict a spectrum of glueballs. The lightest 3 have J^{PC} Quantum numbers of 0^{++} , 2^{++} and 0^{-+} .


The lightest is about $1.6 \text{ GeV}/c^2$





Glueball Mass Spectrum




Morningstar et al.






Glue-rich channels

Where should you look experimentally for Glueballs?




Radiative J/ψ Decays

$0^{-+} \eta(1440)$
 $0^{++} f_0(1710)$
} Large signals



Proton-Antiproton Annihilation



Central Production
(double-pomeron exchange)



Decays of Glueballs?

Glueballs should decay in a flavor-blind fashion.

$$\pi\pi : K\bar{K} : \eta\eta : \eta'\eta' : \eta\eta' = 3 : 4 : 1 : 1 : 0$$

$\eta\eta'=0$ is true for any $SU(3)$ singlet and for any pseudoscalar mixing angle. Only an $SU(3)$ "8" can couple to $\eta\eta'$.

Flavor-blind decays have always been cited as glueball signals. Not necessarily true - coupling proportional to daughter mass can distort this.



Crystal Barrel Results: antiproton-proton annihilation at rest

Discovery of the $f_0(1500)$



Solidified the $f_0(1370)$


Discovery of the $a_0(1450)$

$f_0(1500) \Leftrightarrow \pi\pi, \eta\eta, \eta\eta', KK, 4\pi$

$f_0(1370) \Leftrightarrow 4\pi$

Establishes the scalar nonet





The $f_0(1500)$

Is it possible to describe the $f_0(1500)$ as a member of a meson nonet?


$$\begin{pmatrix} f_0(1370) \\ f_0(1500) \end{pmatrix} = \begin{pmatrix} \cos \theta & \sin \theta \\ -\sin \theta & \cos \theta \end{pmatrix} \begin{pmatrix} 1 \\ 8 \end{pmatrix}$$

Use $SU(3)$ and OZI suppression to compute relative decays to pairs of pseudoscalar mesons

Get an angle of about 143°

90% light-quark

10% strange-quark




Both the $f_0(1370)$ and $f_0(1500)$ are $u\bar{u}$ & $d\bar{d}$



WA102 Results

CERN experiment colliding p on a hydrogen target.

Central Production Experiment



Recent comprehensive data set and a coupled channel analysis.

$$\frac{f_0(1370) \rightarrow \pi\pi}{f_0(1370) \rightarrow K\bar{K}} = 2.17 \pm 0.90$$

$$\frac{f_0(1370) \rightarrow \eta\eta}{f_0(1370) \rightarrow K\bar{K}} = 0.35 \pm 0.21$$

$$\frac{f_0(1500) \rightarrow \pi\pi}{f_0(1500) \rightarrow \eta\eta} = 5.5 \pm 0.84$$

$$\frac{f_0(1500) \rightarrow K\bar{K}}{f_0(1500) \rightarrow \pi\pi} = 0.32 \pm 0.07$$


$$\frac{f_0(1500) \rightarrow \eta\eta'}{f_0(1500) \rightarrow \eta\eta} = 0.52 \pm 0.16$$

$$\frac{f_0(1710) \rightarrow \pi\pi}{f_0(1710) \rightarrow K\bar{K}} = 0.20 \pm 0.03$$

$$\frac{f_0(1710) \rightarrow \eta\eta}{f_0(1710) \rightarrow K\bar{K}} = 0.48 \pm 0.14$$


$$\frac{f_0(1710) \rightarrow \eta\eta'}{f_0(1710) \rightarrow \eta\eta} < 0.05(90\%cl)$$





BES Results

$$J / \psi \rightarrow \gamma X$$






$J / \psi \rightarrow \gamma f_0(1500) \Rightarrow f_0(1500) \rightarrow \pi^+ \pi^-$	$0.665 \cdot 10^{-4}$
$J / \psi \rightarrow \gamma f_0(1500) \Rightarrow f_0(1500) \rightarrow \pi^0 \pi^0$	$0.34 \cdot 10^{-4}$
$J / \psi \rightarrow \gamma f_0(1500) \Rightarrow f_0(1500) \rightarrow \pi^+ \pi^- \pi^+ \pi^-$	$3.1 \cdot 10^{-4}$

$J / \psi \rightarrow \gamma f_0(1710) \Rightarrow f_0(1710) \rightarrow \pi^+ \pi^-$	$2.64 \cdot 10^{-4}$
$J / \psi \rightarrow \gamma f_0(1710) \Rightarrow f_0(1710) \rightarrow \pi^0 \pi^0$	$1.33 \cdot 10^{-4}$
$J / \psi \rightarrow \gamma f_0(1710) \Rightarrow f_0(1710) \rightarrow K \bar{K}$	$9.62 \cdot 10^{-4}$
$J / \psi \rightarrow \gamma f_0(1710) \Rightarrow f_0(1710) \rightarrow \pi^+ \pi^- \pi^+ \pi^-$	$3.1 \cdot 10^{-4}$

Clear Production of $f_0(1500)$ and $f_0(1710)$, no report of the $f_0(1370)$. $f_0(1710)$ has strongest production.




Model for Mixing



$G \rightarrow q\bar{q}$ flavor blind? r
 $u\bar{u}, d\bar{d}, s\bar{s}$

Solve for mixing scheme



Meson Glueball Mixing

Physical Masses

$f_0(1370), f_0(1500), f_0(1710)$

Bare Masses:

m_1, m_2, m_G

$$\frac{(u\bar{u} + d\bar{d})}{\sqrt{2}}$$

$$s\bar{s}$$

	(G)	(S)	(N)
$f_0(1370)$	-0.69 ± 0.07	0.15 ± 0.01	0.70 ± 0.07
$f_0(1500)$	-0.65 ± 0.04	0.33 ± 0.04	-0.70 ± 0.07
$f_0(1710)$	0.39 ± 0.03	0.91 ± 0.02	0.15 ± 0.02

$$\sim (|1\rangle - |G\rangle)$$

$$\sim (|8\rangle - |G\rangle)$$

$$\sim (|1\rangle + |G\rangle)$$

$$m_1 = 1377 \pm 20 \quad m_2 = 1674 \pm 10 \quad m_G = 1443 \pm 24$$

Lattice of about 1600



Glueball Expectations

Antiproton-proton: Couples to $(u\bar{u} + d\bar{d})$

Observe: $f_0(1370), f_0(1500)$

Central Production: Couples to G and $(u\bar{u} + d\bar{d})$ in phase.

Observe: $f_0(1370), f_0(1500)$, weaker $f_0(1710)$.

Radiative J/ψ : Couples to G , $|1\rangle$, suppressed $|8\rangle$

Observe strong $f_0(1710)$ from constructive $|1\rangle+G$

Observe $f_0(1500)$ from G

Observe weak $f_0(1370)$ from destructive $|1\rangle+G$

Two photon: Couples to the quark content of states, not to the glueball. Not clear to me that $\gamma\gamma \rightarrow f_0$ has been seen.



Higher mass glueballs?

Lattice predicts that the 2^{++} and the 0^{-+} are the next two, with masses just above $2\text{GeV}/c^2$.

Radial Excitations of the 2^{++} ground state

$L=3$ 2^{++} States + Radial excitations

$f_2(1950)$, $f_2(2010)$, $f_2(2300)$, $f_2(2340)$...


2'nd Radial Excitations of the η and η' ,
perhaps a bit cleaner environment! (I would
Not count on it though....)

I expect this to be very challenging. Evidence from
BES for an $\eta(1760) \rightarrow \omega\omega$.



Hybrid Mesons

built on quark-model mesons



normal mesons


$$CP = \{(-1)^{L+S}\} \{(-1)^{L+1}\}$$

$$= \{(-1)^{S+1}\}$$

Flux-tube Model

$$m=0 \quad CP = (-1)^{S+1}$$

$$m=1 \quad CP = (-1)^S$$



ρ, ω, ϕ

$$S=0, L=0, m=1$$

$$J=1 \quad CP=+$$

$$J^{PC} = 1^{++}, 1^{--}$$

(not exotic)

$$S=1, L=0, m=1$$

$$J=1 \quad CP=-$$

$$J^{PC} = 0^{-+}, 0^{+-}$$


$$1^{-+}, 1^{+-}$$

$$2^{-+}, 2^{+-}$$

exotic




QCD Potential



Gluonic Excitations provide an experimental measurement of the excited QCD potential.

Observations of exotic quantum number nonets are the best experimental signal of gluonic excitations.



Hybrid Predictions

Flux-tube model: 8 degenerate nonets

$$\underbrace{1^{++}, 1^{--}}_{S=0} \quad \underbrace{0^{-+}, 0^{+-}, 1^{-+}, 1^{+-}, 2^{-+}, 2^{+-}}_{S=1} \quad \sim 1.9 \text{ GeV}/c^2$$

Lattice calculations --- 1^{-+} nonet is the lightest

UKQCD (97)	1.87 ± 0.20
MILC (97)	1.97 ± 0.30
MILC (99)	2.11 ± 0.10
Lacock(99)	1.90 ± 0.20
Mei(02)	2.01 ± 0.10
Bernard(04)	1.792 ± 0.139

1^{-+}	1.9 ± 0.2
2^{+-}	2.0 ± 0.11
0^{+-}	2.3 ± 0.6


In the charmonium sector:

1^{-+}	4.39 ± 0.08	} Splitting = 0.20
0^{+-}	4.61 ± 0.11	



Decays of Hybrids

Decay calculations are model dependent, but the 3P_0 model does a good job of describing normal meson decays.




The angular momentum in the flux tube stays in one of the daughter mesons ($L=1$) and ($L=0$) meson.

$L=0: \pi, \rho, \eta, \omega, \dots$
 $L=1: a, b, h, f, \dots$

} $\eta\pi, \rho\pi, \dots$ not preferred.

$\pi_1 \rightarrow \pi b_1, \pi f_1, \pi \rho, \eta a_1$ 87, 21, 11, 9 MeV (partial widths)
 first lattice prediction ~ 400 MeV





E852 Results

$$\pi^- p \rightarrow \eta \pi^- p$$




(18 GeV)

$$\pi_1(1400) \quad \text{Mass} = 1370 \text{ }^{+16}_{-30} \text{ }^{+50}_{-30} \text{ MeV}/c^2$$

$$\quad \quad \quad \text{Width} = 385 \text{ }^{+65}_{-105} \text{ MeV}/c^2$$

The $a_2(1320)$ is the dominant signal. There is a small (few %) exotic wave.

Interference effects show a resonant structure in 1^{++} . (Assumption of flat background phase as shown as 3.)





Crystal Barrel Results: antiproton-neutron annihilation


$\pi_1(1400)$ Mass = $1400 \pm 20 \pm 20 \text{ MeV}/c^2$
 Width = $310^{+50}_{-30} \text{ MeV}/c^2$

Without π_1 $\chi^2/\text{ndf} = 3$, with = 1.29


Same strength as the a_2 .

Produced from states with **one unit** of angular momentum.





Controversy



In analysis of E852 $\eta\pi^0$ data, so evidence of the $\pi_1(1400)$

In CBAR data, the $\eta\pi^0$ channel is not conclusive.


Analysis by Szczepaniak shows that the exotic wave is not resonant - a rescattering effect.

The signal is far too light to be a hybrid by any model.


This is not a hybrid and may well not be a state.



E852 Results




At 18 GeV/c




$M(\pi^+ \pi^- \pi^-)$ [GeV/c²]



to partial wave analysis



$M(\pi^+ \pi^-)$ [GeV/c²]



suggests





An Exotic Signal

Correlation of
Phase
&
Intensity




Leakage
From
Non-exotic Wave
due to imperfectly
understood acceptance

$$M(\pi^+ \pi^- \pi^-) \text{ [GeV} / c^2 \text{]}$$

$$3\pi \quad m=1593^{+28}_{-47} \quad \Gamma=168^{+150}_{-12}$$

$$\pi\eta' \quad m=1597^{+45}_{-10} \quad \Gamma=340^{+40}_{-50}$$





In Other Channels

E852 Results

$1^- +$ in $\eta'\pi$


$\pi^-p \rightarrow \eta'\pi^-p$ at 18 GeV/c


The $\pi_1(1600)$ is the
Dominant signal in $\eta'\pi$.

Mass = 1.597 ± 0.010 GeV

Width = 0.340 ± 0.040 GeV

$\pi_1(1600) \rightarrow \eta'\pi$








In Other Channels

E852 Results

1-+ in $f_1\pi$ and $b_1\pi$




Mass = 1.709 ± 0.024 GeV

Width = 0.403 ± 0.08 GeV

In both $b_1\pi$ and $f_1\pi$, observe Excess intensity at about $2\text{GeV}/c^2$.

Mass ~ 2.00 GeV,
Width ~ 0.2 to 0.3 GeV



Mass = 1.687 ± 0.011 GeV

Width = 0.206 ± 0.03 GeV



$\pi_1(1600)$ Consistency


3π	$m=1593$	$\Gamma=168$
$\eta'\pi$	$m=1597$	$\Gamma=340$
$f_1\pi$	$m=1709$	$\Gamma=403$
$b_1\pi$	$m=1687$	$\Gamma=206$

**Not Outrageous, but not great agreement.
Mass is slightly low, but not crazy.**

Szczepaniak: Explains much of the η' signal as a background rescattering similar to the $\eta\pi$.



Still room for a narrower exotic state.





New Analysis

Dzierba et. al. PRD 73 (2006)



- Add $\pi_2(1670) \rightarrow \rho\pi$ ($L=3$)
- Add $\pi_2(1670) \rightarrow \rho_3\pi$
- Add $\pi_2(1670) \rightarrow (\pi\pi)_5\pi$
- Add a_3 decays
- Add $a_4(2040)$


10 times statistics in each of two channels.

Get a better description of the data via moments comparison

No Evidence for the $\pi_1(1670)$



Exotic Signals



$\pi_1(1400)$ Width ~ 0.3 GeV, Decays: only $\eta\pi$
 weak signal in πp production (scattering??)
 strong signal in antiproton-deuterium.

NOT A
 HYBRID


$\pi_1(1600)$ Width ~ 0.16 GeV, Decays $\rho\pi, \eta'\pi, (b_1\pi)$
 Only seen in πp production, (E852 + VES)

Does
 this
 exist?


$\pi_1(2000)$ Weak evidence in preferred hybrid
 modes $f_1\pi$ and $b_1\pi$

The right
 place. Needs
 confirmation.





Exotics and QCD




In order to establish the existence of gluonic excitations, We need to establish the existence and nonet nature of the 1^{-+} state.

We need to establish at other exotic QN nonets - the 0^{+-} and 2^{+-} .

In the scalar glueball sector, the decay patterns have provided the most sensitive information. I expect the same will be true in the hybrid sector as well.

DECAY PATTERS ARE CRUCIAL



Summary

The first round of J/ψ experiments opened the door to exotic spectroscopy, but the results were confused.


LEAR at CERN opened the door to precision, high-statistics spectroscopy experiments and significantly improved both our understanding of the scalar mesons and the scalar glueball.

Pion production experiments at BNL (E852) and VES opened the door to states with non-quark-anti-quark quantum numbers. Recent analysis adds to controversy.


CERN central production (WA102) provided solid new data on the scalar sector, and a deeper insight into the scalar glueball.

BES is collecting new J/ψ data. CLEO-c can hopefully add with the ψ' program.







The Future




The GlueX experiment at JLab will be able to do for hybrids what Crystal Barrel and WA102 (together) did for glueballs. What are the properties of static glue in light-quark hadrons and how is this connected to confinement.


The antiproton facility at GSI (HESR) will look for hybrids in the charmonium system - PANDA. May also be able to shed more light on the Glueball spectrum.





Significance of signal.





The Scalar Mesons

Overpopulation
Strange Decay Patterns
Seen in glue-rich reactions
Not in glue-poor

What about 2^{++} and 0^{-+} ?

J/ Ψ Decays?

Awaiting CLEO-c

Glueball and Mesons
are mixed. Scheme is
model dependent.

Crystal Barrel proton-antiproton annihilation






Central Production WA102



Three States

$f_0(1370)$

$f_0(1500)$

$f_0(1710)$





What will we learn?


There may be a signal for hybrid mesons - Exotic quantum numbers - but confirmation requires the observation of a nonet, not just a single state.

Mapping out more than one exotic nonet is necessary to establishing the hybrid nature of the states.

Decay patterns will be useful for the exotic QN states, and necessary for the non-exotic QN states.




Photoproduction of Exotics



Quark spins anti-aligned

A pion or kaon beam,
when scattering occurs,
can have its flux tube excited

Much data in hand with some
evidence for gluonic excitations
(tiny part of cross section)




Quark spins aligned

Almost no data in hand
in the mass region
where we expect to find exotic hybrids
when flux tube is excited





Exotics in Photoproduction




Need to establish nonet nature of exotics: $\pi \eta \eta'$

Need to establish more than one nonet: $0^{+-} 1^{-+} 2^{+-}$





Exotic Hybrid Results




$1^{+-} \pi_1(1400)$

$1^{+-} \pi_1(1600)$

$1^{+-} \pi_1(2000)$





0^{+-} and 2^{+-} Exotics

In photoproduction, couple to ρ , ω or ϕ ?

$$b_0 \quad I^G(J^{PC})=1^+(0^{+-})$$

$$\omega a_1, \rho f_0, \rho f_1$$


$$h_0 \quad I^G(J^{PC})=0^-(0^{+-})$$

$$\omega f_0, \omega f_1, \rho a_1$$

$$h'_0 \quad I^G(J^{PC})=0^-(0^{+-})$$

$$\phi f_0, \phi f_1, \rho a_1$$

$$K_0 \quad I(J^P)=\frac{1}{2}(0^+)$$



$$\omega\pi \quad \omega a_1, \rho f_0, \rho f_1$$

$$b_2 \quad I^G(J^{PC})=1^+(2^{+-})$$

“Similar to π_1 ”

$$\omega\eta, \rho\pi, \omega f_0, \omega f_1, \rho a_1$$

$$h_2 \quad I^G(J^{PC})=0^-(2^{+-})$$



$$\phi\eta, \rho\pi, \phi f_0, \phi f_1, \rho a_1$$

$$h'_2 \quad I^G(J^{PC})=0^-(2^{+-})$$

$$K_2 \quad I(J^P)=\frac{1}{2}(2^+)$$

Kaons do not have exotic QN's







Strong QCD

See $q\bar{q}$ and qqq systems.

Color singlet objects observed in nature:



Nominally, glue is not needed to describe hadrons.



$u \quad \bar{u}$

$d \quad \bar{d}$ Focus on "light-quark mesons"

$s \quad \bar{s}$

Allowed systems: $gg, ggg, q\bar{q}g, q\bar{q}q\bar{q}$

